CONTENTS

	Page.
Relative Water Requirement of Plants. LYMAN J. BRIGGS and H. L. SHANTZ	1
Heart-Rot of Oaks and Poplars Caused by Polyporus Dryophilus. GEORGE G. HEDGCOCK and W. H. LONG	65
Decomposition of Soil Carbonates. W. H. MACINTIRE	79
A Fungous Disease of Hemp. VERA K. CHARLES and ANNA E. JENKINS	81
A More Accurate Method of Comparing First-Generation Maize Hybrids with Their Parents. G. N. COLLINS	85
Natural Revegetation of Range Lands Based upon Growth Requirements and Life History of the Vegetation. ARTHUR W. SAMPSON	93
Pecan Rosette. W. A. ORTON and FREDERICK V. RAND	149
A Nitrogenous Soil Constituent: Tetracarbonimid. EDMUND C. SHOREY and E. H. WALTERS	175
Apple Root Borer. FRED E. BROOKS	179
Changes in Composition of Peel and Pulp of Ripening Bananas. H. C. GORE	187
Assimilation of Colloidal Iron by Rice. P. L. GILE and J. O. CARRERO	205
Coloring Matter of Raw and Cooked Salted Meats. RALPH HOAGLAND	211
Oil Content of Seeds as Affected by the Nutrition of the Plant. W. W. GARNER, H. A. ALLARD, and C. L. FOUBERT	227
Studies in the Expansion of Milk and Cream. H. W. BEARCE	251
Life History of the Melon Fly. E. A. BACK and C. E. PEMBERTON	269
Identification of the Seeds of Species of Agropyron. ROBERT C. DAHLBERG	275
Observations on the Life History of Agrilus Bilineatus. ROYAL N. CHAPMAN	283
Effect of Dilution upon the Infectivity of the Virus of the Mosaic Disease of Tobacco. H. A. ALLARD	295
Moldiness in Butter. CHARLES THOM and R. H. SHAW	301
Susceptibility of Citrous Fruits to the Attack of the Mediterranean Fruit Fly. E. A. BACK and C. E. PEMBERTON	311
Physiological Changes in Sweet Potatoes during Storage. HEINRICH HASSELBRING and LON A. HAWKINS	331
Three-Cornered Alfalfa Hopper. V. L. WILDERMUTH	343
Life History of the Mediterranean Fruit Fly from the Standpoint of Parasite Introduction. E. A. BACK and C. E. PEMBERTON	363

Page.

Relation of Simultaneous Ovulation to the Production of Double-Yolked Eggs. MAYNIE R. CURTIS 375

Brachysm, A Hereditary Deformity of Cotton and Other Plants. O. F. COOK 387

Ability of Colon Bacilli to Survive Pasteurization. S. HENRY AYERS and W. T. JOHNSON, Jr 401

Fitting Logarithmic Curves by the Method of Moments. JOHN RICE MINER 411

Organic Phosphoric Acid of Rice. ALICE R. THOMPSON 425

Two Clover Aphids. EDITH M. PATCH 431

Net Energy Values of Feeding Stuffs for Cattle. HENRY PRENTISS ARMSBY and J. AUGUST FRIES 435

Air and Wind Dissemination of Ascospores of the Chestnut-Blight Fungus. F. D. HEALD, M. W. GARDNER, and R. A. STUDHALTER 493

Index 527

ERRATA

Page 32, Table XX, "*Trifolium repens*" should read "*Trifolium pratense.*"

Page 52, Table XXX, "*Linum usatissimum*" should read "*Linum usitatissimum.*"

Page 52, Table XXX, "*Cucumbis sativa*" should read "*Cucumis sativus.*"

Page 53, Table XXX, "*Boutelona gracilis*" should read "*Bouteloua gracilis.*"

Page 53, Table XXXI, "*Cucumis sativa*" should read "*Cucumis sativus.*"

Page 56, Table XXXIII, column head, "Ratio, 1913 to 1912," should read "Ratio, 1912 to 1913."

Page 56, line 4 from bottom, "75 ± 2" should read "75 ± 1."

Page 97, line 26, "*Salix nutallii*" should read "*Salix nuttallii.*"

Page 106, Table II, "*Sutanion velutinum*" should read "*Sitanion velutinum.*"

Page 165, Table III, insert line after "MgO" to read "Fe_2O_3.... 0.07 0.10 0.09 0.10."

Page 165, Table IV, "Fl_2O_3" should read "Fe_2O_3."

Page 165, Table IV, "Total..... 99.94 ..... 98.36 ....." should read "Total..... 99.94 99.54 100.36 100.00."

Page 166, Table V, "Total..... 104.45 104.82 101.41 101.99" should read "Total ..... 99.73 100.04 99.25 99.81."

Page 256, line 10, "$(C_1 = t - t_m;\ C_2 = C_1^2 - (C_1^2)_m;\ N = D_t - (D_t)_m.)$" should read "$C_1 = t - t_m;\ C_2 = C_1^2 - (C_1^2)_m;\ N = D_t - (D_t)_m$."

Page 256, line 11, "$(\Sigma C_1^2 \alpha + \Sigma C_{12} \beta C = \Sigma C_1 N.)$" should read "$\Sigma C_1^2 \alpha + \Sigma C_1 C_2 \beta = \Sigma C_1 N$."

Page 278, lines 37–38, "(fig. 2)" should read "(fig. 3)."

Page 279, line 19, "as" should read "at."

Page 279, legend for figure 2, "Side views of basal portions, etc.," should read "Fig. 2.—Detail drawings of ventral view of seeds of *Agropyron* spp.: *A*, *Agropyron repens; B*, *A. smithii; C*, *A. tenerum.* × 9."

Page 280, legend for figure 3, "Detail drawings of ventral view, etc.," should read "Fig. 3.—Side views of basal portions of seeds of *Agropyron* spp., showing the relative projection of the rachilla: *A*, *Agropyron repens; B*, *A. smithii; C*, *A. tenerum.* × 9."

Page 280, legend for figure 4, "Edge of rachilla in *Agropyron* spp., etc.," should read "Fig. 4.—Edge of palea in *Agropyron* spp., showing space and comparative size of bristles: *A*, *Agropyron repens; B*, *A. smithii; C*, *A. tenerum.* × 9."

ILLUSTRATIONS

PLATES

Relative Water Requirement of Plants

Page.

Plate I. Fig. 1.—General view of the plant inclosure used at Akron, Colo., showing the pipe framework covered with a hail screen, with the board base surmounted by a single width of cheesecloth to protect the plants against high winds. Fig. 2.—General view inside the inclosure, showing the arrangement of pots and general conditions of growth. Corn and sorghums are shown in the foreground, small grain in the background. Fig. 3.—General view of the inclosure photographed shortly after the grain in some of the pots had been harvested 64

Plate II. Fig. 1.—Pot planted with sugar beets, showing the wax seal around the plants and also the sealed holes where stand was not perfect. Fig. 2.—Weighing pots, showing spring balance, weighing support, and general procedure. Two men operate the weighing support, one of whom lifts the pot by means of a windlass, while a third reads the balance and records the weight. By this method weighings can be made at the rate of two per minute. Fig. 3.—*Grindelia squarrosa* (gumweed) and *Artemisia frigida* (mountain sage), illustrating the growth of native plants used in the water-requirement measurements 64

Plate III. Fig. 1.—Kubanka wheat, grown May 9 to September 3, 1912. Fig. 2.—White Hull-less barley, grown May 16 to August 12, 1912. Fig. 3.—Kubanka wheat. Set grown outside of shelter, May 9 to August 31, 1912. Fig. 4.—Emmer, grown May 11 to August 12, 1912. Fig. 5.—Swedish Select oats, grown May 17 to August 23, 1912. Fig. 6.—Kharkov wheat, grown April 27 to August 28, 1912 64

Plate IV. Fig. 1.—Northwestern Dent corn, grown June 9 to September 16, 1912. Fig. 2.—Hopi corn, grown June 12 to September 26, 1912. Fig. 3.—White durra, grown June 9 to September 26, 1912. Fig. 4.—Red Amber sorghum, grown June 29 to September 27, 1912. Fig. 5.—Minnesota Amber sorghum, grown June 9 to September 26, 1912 64

Plate V. Fig. 1.—Sudan grass. First crop, grown May 28 to July 26, 1912. Fig. 2.—Voronezh proso, grown June 5 to August 20, 1912. Fig. 3.—Kursk millet, grown June 9 to August 20, 1912. Fig. 4.—Select Grimm alfalfa, grown in the open, May 24 to July 27, 1912. Fig. 5.—Select Grimm alfalfa, grown in the shelter, May 24 to July 26, 1912 64

Plate VI. Fig. 1.—Cowpea, grown June 17 to August 26, 1913. Fig. 2.—Hairy vetch, grown May 29 to July 18, 1913. Fig. 3.—Soy bean, grown June 1 to August 26, 1913. Fig. 4.—Cantaloupe, grown June 14 to September 13, 1913. Fig. 5.—Indian Flint corn, grown June 7 to August 27, 1913. Fig. 6.—McCormick potato, grown June 5 to October 4, 1913 64

Plate VII. Fig. 1.—Triumph cotton in shelter, grown May 29 to September 16, 1913. Fig. 2.—*Boebera papposa*, grown July 25 to September 17, 1913. Fig. 3.—Rice in shelter, grown June 12 to September 16, 1913. Fig. 4.—General view of the shelter, showing emmer at the left and White Hull-less barley at the right. Fig. 5.—General view in the shelter, showing corn in the foreground 64

HEART-ROT OF OAKS AND POPLARS CAUSED BY POLYPORUS DRYOPHILUS

Page.

PLATE VIII. Fig. 1.—*Quercus alba:* Crescent-shaped "soak," the initial stage of the piped rot produced by *Polyporus dryophilus;* from Arkansas. Fig. 2.—*Quercus alba:* A radial view of the rot in a limb, showing delignification; from Arkansas. Fig. 3.—*Quercus oblongifolia:* A radial view of rot, showing delignification; from Arizona. Fig. 4.—*Quercus alba:* A final stage of the rot, radial view, with more complete delignification; from Arkansas. Fig. 5.—*Quercus alba:* A tangential view of the rot, showing delignification in pockets; from Arkansas. Fig. 6.—*Quercus alba:* An end view showing a cross section from the same tree as the preceding; from Arkansas. Fig. 7.—*Quercus* sp.: A section of oak from Von Tubeuf, sent to the junior writer as a specimen of the rot caused by *Polyporus dryadeus* in Europe. Fig. 8.—*Quercus* sp.: The reverse side of the specimen shown in the preceding. Fig. 9.—*Quercus* sp.: A section of oak from Europe, obtained by Von Schrenk, with a piped rot similar to that of *Polyporus dryophilus*.... 78

PLATE IX. Fig. 1.—A sporophore of *Polyporus dryophilus,* tuberous form on *Quercus gambelii;* from Arizona. Fig. 2.—Sectional view of a sporophore of *Polyporus dryophilus* on *Quercus gambelii,* showing the hard granular core with whitish mycelial strands; also the pore layer; from New Mexico. Fig. 3.—A sporophore of *Polyporus dryophilus* on *Quercus californica,* showing the upper surface with a faint zonation; from California. Fig. 4.—A section through a sporophore of *Polyporus dryophilus* on *Quercus garryana,* showing the structure of the hard granular core; from California. Fig. 5.—A front view, showing the margin of the same sporophore as in figure 3, representing the ungulate form. Fig. 6.—A view of the pore surface of an applanate sporophore of *Polyporus dryophilus* on *Quercus alba;* from Arkansas 78

PLATE X. Fig. 1.—A sporophore of *Polyporus dryophilus,* front view showing the margin, on *Populus tremuloides;* from Colorado. Fig. 2.—A second sporophore from the same tree as figure 1, showing an imbricated form. Fig. 3.—A view of the upper surface of a sporophore of *Polyporus rheades* on *Populus tremula;* from Stockholm, Sweden. Fig. 4.—A sectional view of a sporophore of *Polyporus corruscans* on Quercus; from Upsala, Sweden. Fig. 5.—A side view of an imbricate sporophore of *Polyporus dryophilus,* applanate form on *Populus tremuloides;* from Colorado. Fig. 6.—A sectional view of the same sporophore as in the preceding figure, showing the hard granular core and whitish mycelial strands. Fig. 7.—A view of the upper surface of an applanate sporophore of *Polyporus dryophilus* on *Quercus alba;* from Arkansas. Fig. 8.—The pore surface of a sporophore of *Polyporus dryophilus* on *Populus tremuloides;* from Colorado..... 78

A FUNGOUS DISEASE OF HEMP

PLATE XI. A hemp plant, showing upper branches attacked by the fungous *Botryosphaeria marconii* 84

NATURAL REVEGETATION OF RANGE LANDS BASED UPON GROWTH REQUIREMENTS AND LIFE HISTORY OF THE VEGETATION

PLATE XII. Fig. 1.—View of the lower grazing lands in the Wallowa National Forest. Fig. 2.—Characteristic open stand of western yellow pine and dense cover of herbaceous vegetation, mainly pine-grass (*Calamagrostis pubescens*), Wallowa National Forest. Transition zone (yellow-pine association). Fig. 3.—A burned-over area of lodgepole pine, with characteristic dense sapling stand..... 148

Page.

PLATE XIII. Fig. 1.—Dense stand of lodgepole pine, with undergrowth of red huckleberry (*Vaccinium scoparium*). Canadian zone (lodgepole-pine association). Fig. 2.—A flat eminence in the Hudsonian zone, showing the characteristic clumped growth of whitebark pine and Alpine fir. Fig. 3.—Irregular topography of the upper grazing lands. Hudsonian zone (whitebark-pine association) 148

PLATE XIV. Fig. 1.—Arctic-Alpine and upper-subalpine region, where forage is sparse, due to poor soil, short growing season, and unfavorable climate. Fig. 2.—Mountain range lands prior to the beginning of growth and germination. Fig. 3.—Same view as shown in figure 2, but more in detail, showing the condition eight days later (June 30) 148

PLATE XV. Fig. 1.—Contrast in the progress of the flower stalk production of mountain bunch-grass on portion of range which has been completely closed to grazing for a period of three successive years and on range which has been subject to continued early grazing. Fig. 2.—Western porcupine grass (*Stipa occidentalis*), showing empty glumes and floret with the scale and its awned projection to the left; to the right the floret with glumes removed, showing the sharp-pointed, slightly curved seed tip. Fig. 3.—Average development of the root system and aerial portion of mountain bunch-grass at end of the first growing season 148

PLATE XVI. Mountain bunch-grass, showing root development and aerial growth at the end of the second season 148

PLATE XVII. Mountain bunch-grass in the spring of the third year of growth just before producing flower stalks, showing the natural position and length of the elaborate root development and aerial growth 148

PLATE XVIII. Mountain bunch-grass at the end of the third year, showing three flower stalks and inflorescence 148

PLATE XIX. Sickle sedge (*Carex umbellata brevirostris*), showing offshoots from the rootstocks and flower stalks with fruit in the process of development. 148

PLATE XX. Fig. 1.—Station 4 on Stanley Range as it appeared on July 12, 1907. Fig. 2.—View of station 4 on July 15, 1909, after two years' protection from grazing animals. Fig. 3.—View of quadrat 1, established on July 10, 1907. 148

PLATE XXI. Fig. 1.—Quadrat 1, as it appeared on July 16, 1909. Fig. 2.—Area of mountain bunch-grass closed to grazing animals on July 8, 1907. Fig. 3.—View of open range contiguous to area shown in figure 2 148

PLATE XXII. View of plot in the Transition (yellow-pine) zone which has been protected from grazing animals for three successive years, showing contrast in carrying capacity with contiguous open range 148

PLATE XXIII. Fig. 1.—View of portion of allotment at medium elevation where the destruction of forage seedlings due to grazing and trampling was studied. Fig. 2.—Dense stand of smooth wild rye (*Elymus glaucus*) and short-awned brome-grass (*Bromus marginatus*) seedlings 148

PECAN ROSETTE

PLATE XXIV. Fig. 1.—One normal pecan leaf and two leaves with rosette from Dewitt, Ga. Fig. 2.—Pecan shoot with early symptoms of rosette.... 174

PLATE XXV. Rosetted pecan leaf showing perforations due to the failure of part of the mesophyll to develop 174

PLATE XXVI. Fig. 1.—Pecan shoot in advanced stages of rosette. Fig. 2.—Normal pecan shoot for comparison with rosetted shoot 174

PLATE XXVII. Fig. 1.—Young orchard pecan tree with a moderate attack of rosette on the left side and seriously dying back from the disease on the other side. Fig. 2.—Young orchard pecan tree in advanced stages of rosette 174

Page.

PLATE XXVIII. Fig. 1.—Young orchard tree with severe attack of rosette. Fig. 2.—Rosetted pecan tree cut off to the stump the preceding season, with the present season's growth again distinctly showing rosette. Fig. 3.—Two seedling pecan trees planted the same day from the same lot of seedlings 174

APPLE ROOT BORER

PLATE XXIX. Figs. 1 and 3.—Sections of an apple root, showing burrows of the apple root borer (*Agrilus vittaticollis*). Fig. 2.—Cross section of the trunk of a young apple tree, showing burrows made by the larvæ of the apple root borer in ascending the trunk to pupate 186

PLATE XXX. Fig. 1.—*Agrilus vittaticollis:* Larva (*c*), pupa (*b*), and adult (*a*) of the apple root borer in the pupal cell. Fig. 2.—*Xylophruridea agrili*, a common parasite of the apple root borer. Fig. 3.—Section of trunk of young service tree, showing below the white egg and above the exit hole of the apple root borer. Fig. 4.—*Xylophruridea agrili:* Larvæ of the parasite; one feeding on the larva and the other in the pupal cell of its host ... 186

PLATE XXXI. Fig. 1.—*Agrilus vittaticollis:* Egg on trunk of young service tree. Fig. 2.—*Agrilus vittaticollis:* Feeding form of larva. Fig. 3.—*Agrilus vittaticollis:* Contracted form of larva as taken from pupal cell. Fig. 4.—*Agrilus vittaticollis:* Pupa. Fig. 5.—*Agrilus vittaticollis: a*, Adult, or beetle; *b*, claw; *c*, antenna. Fig. 6.—*Xylophruridea agrili*, a parasite of the apple root borer: Pupa 186

COLORING MATTER OF RAW AND COOKED SALTED MEATS

PLATE XXXII. Fig. 1.—Oxyhemoglobin, ox blood. Fig. 2.—Oxyhemoglobin, ox blood. Fig. 3.—NO-hemoglobin, ox blood. Fig. 4.—Methemoglobin, ox blood. Fig. 5.—Methemoglobin, ox blood 226

PLATE XXXIII. Fig. 1.—Oxyhemoglobin, sheep blood. Fig. 2.—Oxyhemoglobin, sheep blood. Fig. 3.—NO-hemoglobin, sheep blood. Fig. 4.—NO-hemoglobin, pig blood. Fig. 5.—NO-hemoglobin, pig blood 226

IDENTIFICATION OF THE SEEDS OF SPECIES OF AGROPYRON

PLATE XXXIV. *Agropyron repens:* Spikes showing degrees of variation which may occur. *A*, Typical spike 282

PLATE XXXV. *Agropyron smithii:* Spikes showing degrees of variation. *A*, Typical spike 282

PLATE XXXVI. *Agropyron tenerum:* Spikes showing degrees of variation. *A*, Typical spike 282

PLATE XXXVII. *Agropyron* spp.: Typical seeds and spikelets. Fig. 1.—*Agropyron repens.* Fig. 2.—*Agropyron smithii.* Fig. 3.—*Agropyron tenerum* .. 282

OBSERVATIONS ON THE LIFE HISTORY OF AGRILUS BILINEATUS

PLATE XXXVIII. Fig. 1.—*Agrilus bilineatus:* Eggs in position in the bark of an oak tree. Fig. 2.—*Agrilus bilineatus:* Cluster of newly laid eggs. Fig. 3.—*Agrilus bilineatus:* Eggs shortly before hatching. Fig. 4.—*Agrilus bilineatus:* Newly hatched larva. Fig. 5.—*Agrilus bilineatus:* Mature larva. Fig. 6.—*Agrilus bilineatus:* Larva in its cell. Section made perpendicular to the surface of the bark. *A*, Point at which adult will emerge; *B*, burrow stopped with frass. Fig. 7.—*Agrilus bilineatus:* Pupa in cell. Section made parallel to the surface of the bark. Fig. 8.—*Agrilus bilineatus:* Adult female. Fig. 9.—*Agrilus bilineatus:* Adult male 294

Page.

PLATE XXXIX. Fig. 1.—Leaf showing work of four Agrilus beetles in 24 hours. Fig. 2.—Hole in bark made by adult Agrilus in emerging from pupal cell. Fig. 3.—Larvæ of *Agrilus bilineatus* and their burrows. Fig. 4.—Complete burrow of a larva of *Agrilus bilineatus*. *A*, Point at which larva hatched; *B*, beginning of second instar; *C*, beginning of third instar; *D*, beginning of fourth instar; *E*, pupal cell 294

SUSCEPTIBILITY OF CITROUS FRUITS TO THE ATTACK OF THE MEDITERRANEAN FRUIT FLY

PLATE XL. Fig. 1.—Orange infested with larvæ of the Mediterranean fruit fly (*Ceratitis capitata*). Fig. 2.—Orange infested with larvæ of the Mediterranean fruit fly (*Ceratitis capitata*), showing two breathing holes of the larvæ in the decayed area 330

PLATE XLI. Cross section of shaddock No. 1, showing the thick, loose texture of the rag with darkened area above and to the right showing the channels made by well-grown Mediterranean fruit-fly larvæ 330

PLATE XLII. Fig. 1.—Cross section of the orange shown on Plate XL, figure 2. Fig. 2.—Orange containing 87 punctures in the rind 330

THREE-CORNERED ALFALFA HOPPER

PLATE XLIII. Fig. 1.—The three-cornered alfalfa hopper (*Stictocephala festina*): Adult. *a*, View from side; *b*, view from front. Fig. 2.—The three-cornered alfalfa hopper: *a*, Nymph in first stage; *b*, egg. Fig. 3.—The three-cornered alfalfa hopper: Nymph in second stage. Fig. 4.—The three-cornered alfalfa hopper: Nymph in third stage. Fig. 5.—The three-cornered alfalfa hopper: Nymph in fourth stage. Fig. 6.—The three-cornered alfalfa hopper: Nymph in fifth stage. Fig. 7.—An alfalfa stem showing feeding punctures of the three-cornered alfalfa hopper: *a*, Ring or girdle of punctures around the stem; *b*, gall resulting from girdling 362

LIFE HISTORY OF THE MEDITERRANEAN FRUIT FLY FROM THE STANDPOINT OF PARASITE INTRODUCTION

PLATE XLIV. Fig. 1.—Wooden boxes, 14 by 12 by 3 inches in size, used in obtaining pupæ of fruit flies. Fig. 2.—Contrivance used for keeping the infested fruit free from the sand and bringing the emerging larvæ to a central container where they may be gathered quickly 374

PLATE XLV. Fig. 1.—Method of keeping adult fruit flies alive over long periods. Fig. 2.—An apple after having been suspended for one day in a jar containing Mediterranean fruit flies 374

RELATION OF SIMULTANEOUS OVULATION TO THE PRODUCTION OF DOUBLE-YOLKED EGGS

PLATE XLVI.—Fig. 1.—Large yolk (weight, 30.12 gm.) with two germ disks; found in a large hen's egg. Fig. 2.—Fused immature yolks (weight, 1.45 gm.); found in a small hen's egg. Fig. 3.—Type I double-yolked egg, showing two yolks with separate vitelline membranes but inclosed in a common chalaziferous layer 386

PLATE XLVII. Fig. 1.—Type I double-yolked egg, showing two yolks with separate vitelline membranes but inclosed in a common chalaziferous layer. Fig. 2.—Type II double-yolked egg, showing two yolks with separate chalazal membranes but common thick albumen 386

Page.

PLATE XLVIII. Fig. 1.—Type II double-yolked egg, showing two yolks with some separate and some common thick albumen envelopes. Fig. 2.—Type III double-yolked egg, showing two yolks with all the thick albumen separate 386

PLATE XLIX. Fig. 1.—Shell of type III double-yolked egg, which shows external evidence of its double nature by a seam in the shell. Fig. 2.—The inside of the shell shown in figure 1, showing the fold of egg membrane which projected between the two component eggs 386

PLATE L. Oviduct removed from a laying bird and cut open along the point of attachment of the ventral ligament. *A*, Funnel; *B*, albumen-secreting region; *X*, isthmus ring; *C*, isthmus; *D*, shell gland; and *E*, vagina 386

PLATE LI. Fig. 1.—Ovary of a pullet, showing the follicles which produced the yolks for the double-yolked egg shown in Plate XLVIII, figure 1. Fig. 2.—Ovary of a pullet, showing follicles which produced the yolks for a double-yolked egg similar in structure to the one shown in Plate XLVIII, figure 2 386

PLATE LII. Fig. 1.—Ovary of a pullet, showing a series of resorbing follicles, two of which (probably *C* and *C'*) produced the yolks for the double-yolked egg shown in Plate XLVI, figure 3. Fig. 2.—Ovary of a bird, showing the two largest resorbing follicles, one of which produced the yolk with two germ disks shown in Plate XLVI, figure 1 386

BRACHYSM, A HEREDITARY DEFORMITY OF COTTON AND OTHER PLANTS

PLATE LIII. Abnormal simple leaf on fruiting branch of Egyptian cotton, accompanied by abnormal leaf-like bract, remainder of involucre and floral bud removed 400

PLATE LIV. Normal 3-lobed leaf of fruiting branch of Egyptian cotton, accompanied by normal involucral bract for comparison with Plate LIII 400

PLATE LV. Abnormal leaf of fruiting branch of Egyptian cotton with one stipule enlarged and the lobe of the same side wanting 400

PLATE LVI. Brachytic fruiting branches of "cluster" cotton (Willets Red Leaf) shortened to a single internode by abortion of terminal bud. Fig. 1.—The boll at the right is borne by a very short branch from an axillary bud. Fig. 2.—The boll at the right is borne by the shortened fruiting branch. The left-hand boll represents a shortened branch in the axil of the leaf that subtends the fruiting branch 400

PLATE LVII. Normal and brachytic joints on same fruiting branch of Upland cotton 400

PLATE LVIII. Branches of abnormal variation of Upland cotton, with abortive buds remaining attached to branches by decurrent pedicels and elongated bud scars. The left-hand branch shows abnormal inequality in the lengths of the internodes 400

PLATE LIX. Portion of brachytic fruiting branch of Simpkins cotton producing twin fasciated branches from an axillary bud 400

PLATE LX. Portion of fruiting branch of Columbia cotton, with one internode adnate to the pedicel of the boll of the preceding internode 400

PLATE LXI. Fig. 1.—Plant of Dale Egyptian cotton, showing complete abortion of fruiting branches on the main stalk, while the vegetative branches of the same plant produced a few fruiting branches and ripened a few bolls. Fig. 2.—End of main stalk of plant shown in figure 1, showing abortion of terminal bud and compensatory thickening of the petioles 400

Page.

PLATE LXII. Ends of main stalks of two plants of Dale Egyptian cotton, showing simple fruiting branches and closely similar axillary fruiting branches. 400

AIR AND WIND DISSEMINATION OF ASCOSPORES OF THE CHESTNUT-BLIGHT FUNGUS

PLATE LXIII. Fig. 1.—Petri-dish culture 5044 from 12 minutes' exposure of chestnut-bark agar, made on September 20, 1913, 2 hours and 8 minutes after the cessation of a rain, at station 51, located 27 feet from the nearest lesion. Fig. 2.—Petri-dish culture 5041 from 16 minutes' exposure of chestnut-bark agar, made on September 20, 1913, 1 hour and 55 minutes after the cessation of a rain, at station 49, located 414 feet from the source of the spores. 526

PLATE LXIV. Fig. 1.—Ascospore trap 51. This consists of a wooden bracket which supports an object slide over perithecial pustules. Fig. 2.—Ascospore trap 52. Fig. 3.—Water spore trap located at Station V. 526

PLATE LXV. Fig. 1.—View looking toward the coppice growth from water spore-trap Station V. Fig. 2.—View of a mixed chestnut and oak grove taken from water spore-trap Station VI. 526

TEXT FIGURES

RELATIVE WATER REQUIREMENT OF PLANTS

FIG. 1. Evaporation from a free-water surface (tank) at Akron, Colo., in 1911 and 1912. 8

A FUNGOUS DISEASE OF HEMP

FIG. 1. Microscopic characters of the hemp fungus *Botryosphaeria marconii*. *A*, Sketch of a section of stroma from culture, showing developing perithecia: *a*, microconidial stage, *b*, ascosporic stage. *B*, An ascus with ascospores. *C*, Ascospores. *D*, Macroconidia. *E*, Conidiophores of the Dendrophoma stage. *F*, Microconidia 82

NATURAL REVEGETATION OF RANGE LANDS BASED UPON GROWTH REQUIREMENTS AND LIFE HISTORY OF THE VEGETATION

FIG. 1. Curve showing the variation in the mean temperature in the Transition, Canadian, and Hudsonian grazing zones in 1909. 98

2. Diagram showing the total precipitation in the Transition, Canadian, and Hudsonian grazing zones during July, August, and September, 1909, inclusive. 99

3. Curve showing the comparative daily evaporation in the Transition, Canadian, and Hudsonian zones in 1909. 100

4. Curve showing the maximum and minimum temperature records in the Hudsonian zone (whitebark-pine association). 110

5. Chart of permanent and denuded quadrats 1 and 2 in station 4, established on July 3, 1907. 122

6. Chart of permanent and denuded quadrats 1 and 2 in station 4, remapped on July 12, 1909. 123

PECAN ROSETTE

FIG. 1. Map showing the known distribution of pecan rosette in the United States. 149

CHANGES IN COMPOSITION OF PEEL AND PULP OF RIPENING BANANAS

Page.

FIG. 1. Constant-temperature humidor 193

COLORING MATTER OF RAW AND COOKED SALTED MEATS

FIG. 1. Spectra of hemoglobin and some of its derivatives: *A*, Absorption spectrum of a solution of oxyhemoglobin; *B*, absorption spectrum of a solution of NO-hemoglobin; *C*, absorption spectrum of a solution of hemoglobin prepared by treating a solution of oxyhemoglobin with hydrazin hydrate; *D*, absorption spectrum of a solution of methemoglobin prepared by treating a solution of oxyhemoglobin with potassium ferricyanid; *E*, absorption spectrum of an alkaline solution of hematin; *F*, absorption spectrum of a solution of hemochromogen prepared by treating an alkaline solution of hematin with hydrazin hydrate; *G*, absorption spectrum of a solution of NO-hemochromogen 215

STUDIES IN THE EXPANSION OF MILK AND CREAM

FIG. 1. Specific gravity of milk and cream at 35°/4° C., showing value of α and β 261

2. Specific gravity of milk and cream at 35°/4° C., showing relation between density and percentage of butter fat 261

IDENTIFICATION OF THE SEEDS OF SPECIES OF AGROPYRON

FIG. 1. Detail drawings of dorsal view of *Agropyron* spp.: *A*, *Agropyron repens; B*, *A. smithii; C*, *A. tenerum* 278

2. Detail drawing of ventral view of seeds of *Agropyron* spp.: *A*, *Agropyron repens; B*, *A. smithii; C*, *A. tenerum*. (See "Errata.") 279

3. Side views of basal portions of seeds of *Agropyron* spp., showing the relative projection of the rachilla: *A*, *Agropyron repens; B*, *A. smithii; C*, *A. tenerum*. (See "Errata.") 280

4. Edge of rachilla in *Agropyron* spp., showing shape and comparative size of bristles: *A*, *Agropyron repens; B*, *A. smithii; C*, *A. tenerum* 280

MOLDINESS IN BUTTER

FIG. 1. Graph showing the effect of salt on molding 308

SUSCEPTIBILITY OF CITROUS FRUITS TO THE ATTACK OF THE MEDITERRANEAN FRUIT FLY

FIG. 1. Cross section of peach, showing egg cavity of the Mediterranean fruit fly with eggs 320

2. Cross section of peach, showing the general shriveling of the walls of the egg cavity and the separation of the eggs 320

3. Section of grapefruit rind, showing two egg cavities, one in cross section. 321

THREE-CORNERED ALFALFA HOPPER

FIG. 2. Map showing distribution of the three-cornered alfalfa hopper (*Stictocephala festina*) in the United States 344

ABILITY OF COLON BACILLI TO SURVIVE PASTEURIZATION

FIG. 1. Curve showing results of heating cultures of colon bacilli for 30 minutes at various temperatures 404

TWO CLOVER APHIDS

Page.
FIG. 1. *Aphis brevis:* Antenna of fall alate female collected from hawthorn.... 432
2. *Aphis brevis:* Antenna of alate male..... 433
3. *Aphis bakeri:* Antenna of alate female collected from clover..... 433

NET ENERGY VALUES OF FEEDING STUFFS FOR CATTLE

FIG. 1. Graph showing the dry matter eaten and the increments of heat production due to standing, computed per 500 kg. live weight per 24 hours. 455
2. Graph showing the relation of heat production to dry matter consumed, computed per 500 kg. live weight..... 472

AIR AND WIND DISSEMINATION OF ASCOSPORES OF THE CHESTNUT-BLIGHT FUNGUS

FIG. 1. Map of chestnut coppice growth at West Chester, Pa., in and near which the experiments on wind dissemination of the chestnut-blight fungus were carried out..... 497
2. Map showing the location of some of the important outlying exposure-plate stations..... 498
3. Map showing the location of water spore-trap stations Nos. I to VI.... 520

ADDITIONAL COPIES

OF THIS PUBLICATION MAY BE PROCURED FROM
THE SUPERINTENDENT OF DOCUMENTS
GOVERNMENT PRINTING OFFICE
WASHINGTON, D. C.
AT
25 CENTS PER COPY

SUBSCRIPTION PER YEAR, 12 NUMBERS, $2.50

V31
001
14
XUM